

**BASIC FIBROBLAST GROWTH FACTOR (bFGF) AND
FIBROBLAST GROWTH FACTOR RECEPTOR 1 (FGFR1)
APPEARANCE IN CONGENITAL INTRA-ABDOMINAL ADHESIONS
IN CHILDREN UNDER ONE YEAR OF AGE**

Author: Anna Augule¹

Supervisor: Māra Pilmāne²

¹ Rīga Stradiņš University, Faculty of Medicine, Latvia

² Rīga Stradiņš University, Institute of Anatomy and Anthropology, Latvia

Key words: intra-abdominal adhesions, growth factors.

Introduction. Congenital intra-abdominal adhesions occur due to disruption of the normal embryologic development. Interruption of typical intestinal rotation and fixation during fetal development can occur at a wide range of locations. The most common type found in paediatric patients is incomplete rotation predisposing to midgut volvulus. There are limited publications that investigate the possible aetiology and pathogenesis of congenital peritoneal adhesions.

The aim of this work was to investigate expression of bFGF and FGFR1 in congenital intra-abdominal adhesions.

Material and Methods. The specimens used for research were obtained from 50 patients aged 1 to 292 days who underwent abdominal surgery due to the obstructive gut malrotation and several additional pathologies. Specimens are property of the collection of the Institute for Anatomy and Anthropology of Riga Stradiņš University. Tissues were processed for bFGF and FGFR1 by means of biotin-streptavidin immunohistochemistry.

For quantification of structures, the semiquantitative counting method was used. The designations were as follows: 0, negative reaction; 0/+, occasionally marked structures in the view field; +, a few positive structures in the view field; +/++ few to moderate positive structures; ++, a moderate number of marked structures in the view field; ++/+++ moderate to numerous number of positive structures; +++, a numerous number of marked structures in the view field; ++++, abundance of marked structures found in the view

field. Data analysis was conducted using Statistical Package for the Social Sciences (SPSS) program version 20.0.

Results. bFGF was seen exclusively in fibroblasts and macrophages. In four cases the number of marked cells was abundant, in five cases numerous, but in two cases moderate to numerous. Eleven specimens showed a moderate number of bFGF positive cells. Two specimens showed few to moderate bFGF positive cells, 15 – few positive cells. Occasional positive fibroblasts and macrophages were observed in four cases. 15 specimens showed a completely negative reaction to bFGF.

A numerous number of FGFR1 fibroblasts and macrophages was observed five cases. In another eight cases a moderate to numerous number positive structures was marked. FGFR1 positive structures were mostly seen in moderate (13 cases) and few to moderate (12 cases) appearance. Few fibroblasts and macrophages contained this factor in nine specimens. Occasional positive structures were observed in three cases.

Using the Pearson's correlation test a strong positive correlation was observed between the immunoreactive structures for bFGF and FGFR1 ($r = 0.523$; $p < 0,001$).

Conclusions. Connection between the less distinct bFGF and more prominent FGFR1 proves the compensatory stimulation of receptors as a response on the lack of the same factor in course of adhesion disease.