

Arts therapies

What is the evidence that arts therapies can reduce depression and increase wellbeing across the life span? Relevant examples and important considerations from process and outcome studies

Prof Vicky Karkou

Edge Hill University

Research Centre for Arts and Wellbeing







Research team and scope




























Aim of presentation



- Arts Therapies: What are they?
- Outcome studies: Do they work?
 With whom?
- Therapeutic factors: What supports therapeutic change?
- Future research: What next?






Definition

Arts therapies (or arts psychotherapies) are:

"... the creative use of the artistic media as vehicles for non-verbal and/or symbolic communication, within a holding environment, encouraged by a well-defined client-therapist relationship, in order to achieve personal and/or social therapeutic goals appropriate for the individual" (Karkou and Sanderson 2006, p. 46).

Four sub-disciplines in the UK:

Art, music, drama and dance movement therapy





Professional status in the UK

Training

2-year-long Masters (MA or MSc) University-based qualifications


Registration

- Health and Care Professions Council, next to clinical psychologists, counselling psychologists, occupational therapists and physiotherapists
- UK Council for Psychotherapy, next to psychoanalysts and humanistic psychotherapists
- British Association for Counselling and Psychotherapy next to personcentred counsellors and cognitive behavioural therapists




Areas of work

Main work environments



Main client groups






Working with children






Children with cancer, asthma, those born preterm, ADHD, autism






Children and young people



First published: 01 March 2023 | https://doi.org/10.1002/jcv2.12145 | Citations: 2






Children and Youth Services Review
Volume 112, May 2020, 104928



A systematic review of child-focused outcomes and assessments of arts therapies delivered in primary mainstream schools





Although many practitioners work with children and young people, outcome studies in this area remain limited.



Examples from research in schools







Children with emotional/behavioural difficulties

Improvements sustained for 12 months after the intervention:

- Quality of Life self reported (HRQOL)
- Wellbeing and life functioning self reported (CORS)
- Emotional and behavioural difficulties assess by teachers (SDQ)
- Quality of sleep (fitbits)
- No differences between different types of arts therapies (Moula, Powell and Karkou 2020)

















More outcome studies

Schools

Feasibility RCT in 5 schools in the North West of England with a trauma-informed creative therapy groups.



Hospitals

Feasibility RCT in specialist mental health services in hospitals as a group treatment for children and young people with symptoms of depression.

National Institute for Health Research



Working with adults





Adults with PTSD, anxiety, cancer, heart conditions, pain


Art Therapy and Cognitive Processing Therapy for Combat-Related PTSD: A Randomized Controlled

WILEY

Melissa Campbell, Kathleen P. Decker , Kerry Kruk & Sarah P. Deaver Pages 169-177 | Published online: 20 Oct 2016



Check for updates



A randomized, controlled trial of mindfulness-based art therapy (MBAT) for women with cancer

Daniel A. Monti 🔀, Caroline Peterson, Elisabeth J. Shakin Kunkel, Walter W. Hauck, Lora Rhodes, George C. Brainard

First published: 15 November 2005 | https://doi.org/10.1002/pon.988 | Citations



The Arts in Psychotherapy Volume 39, Issue 4, September 2012, Pages 296-303

The efficacy of dance movement therapy group on improvement of quality of life: A randomized controlled trial

Iris Bräuninger PhD, BTD 🙏 🖾

+ Add to Mendeley & Share 55 Cite



Effect of Music Therapy with Emotional-Approach Coping on Preprocedural Anxiety in Cardiac Catheterization: A Randomized Controlled Trial

Claire M. Ghetti, PhD, LCAT, MT-BC 🗷

Journal of Music Therapy, Volume 50, Issue 2, Summer 2013, Pages 93–122, https://doi.org/10.1093/jmt/50.2.93 Published: 01 July 2013

CLINICAL TRIAL article

Front. Psychol., 16 April 2021

Sec. Psychology for Clinical Settings Volume 12 - 2021 |

https://doi.org/10.3389/fpsyg.2021.587923

This article is part of the Research Topic The Psychological and Physiological Benefits of the Arts

View all 84 articles >

The Development of Mindful-Based Dance Movement Therapy Intervention for Chronic Pain: A Pilot Study With Chronic Headache









Adults



Cochrane Database of Systematic Reviews

Music therapy for depression (Review)

Aalbers S, Fusar-Poli L, Freeman RE, Spreen M, Ket JCF, Vink AC, Maratos A, Crawford M, Chen XJ,



Trusted evidence. Informed decisions.

Cochrane Reviews

Trials -

Clinical Answers <

About ▼

Help ▼

Cochrane Database of Systematic Reviews

Dance movement therapy for depression



Complementary Therapies in Clinical Practice Volume 41, November 2020, 101246



Effectiveness of mindfulness-based art therapy for symptoms of anxiety, depression, and fatigue: A systematic review and meta-analysis

Pamela Newland ^a A

■ B. Ann Bettencourt ^b Show more V + Add to Mendeley < Share 55 Cite https://doi.org/10.1016/j.ctcp.2020.101246 Get rights and conter Systematic reviews of studies with adults with depression

SYSTEMATIC REVIEW article

Front. Psychol., 03 May 2019 Sec. Psychology for Clinical Settings

Volume 10 - 2019 https://doi.org/10.3389/fpsyg.2019.00936

This article is part of the Research Topic

The State of the Art in Creative Arts

Effectiveness of Dance Movement Therapy in the Treatment of Adults With Depression: A





ARTS THERAPIES IN THE TREATMENT OF DEPRESSION

Ania Zubala and Vicky Karkou



There is a growing number of studies that testify on the effectiveness of arts therapies on measures of depression. However, most studies do not have an active control.



Examples from research in the NHS

ERA study – a multi-site trial on group arts therapies with people with diagnostically heterogeneous psychiatric patients; NIHR - HTA



The Arts in Psychotherapy

Volume 72, February 2021, 101747



Research Article

What makes us the same? What makes us different? Development of a shared model and manual of group therapy practice across art therapy, dance movement therapy and music therapy within community mental health care

Catherine Carr ^{a b} ♀ ☒, Barbara Feldtkeller ^c ☒, Jennifer French ^b ☒, Dominik Havsteen-Franklin $d \in \mathbb{Z}$, Val Huet $f \subseteq \mathbb{Z}$, Vicky Karkou $g \subseteq \mathbb{Z}$, Stefan Priebe $g \in \mathbb{Z}$ Stephen Sandford b

counselling & psychotherapy research Linking Research with Practice





Bringing creative psychotherapies to primary NHS Mental Health Services in the UK: A feasibility study on patient and staff experiences of arts for the blues workshops delivered at Improving Access to Psychological Therapies (IAPT) services

Vicky Karkou X, Joanna Omylinska-Thurston, Ailsa Parsons, Kerry Nair, Jennifer Starkey, Shelly Haslam, Scott Thurston, Linda Dubrow Marshall

First published: 02 June 2022 | https://doi.org/10.1002/capr.12544 | Citations: 4





Arts for the Blues

What is it?
What do participants think?



Who is this group for?

Adults:

- Primary mental health care GMMH NHS FT
- University of Salford Clinic
- Mothers and babies /
 parents and infants /
 mothers and pre-school
 children with lived
 experience of domestic
 abuse community
 organisations

Children and young people:

- Schools in the North West of England
- Children's hospital Alder Hey Children's NHS FT CAMHS
- Training for therapists -Greater Manchester CAMHS
- RCT feasibility study RfPB application

Helping professionals:


- Multi-level intervention via Staff Psychology Support Service at large NHS Trust - LUFT
- Doctors in training HEE funding
- Mental health staff -RfPB application in preparation

Working with older adults





Older adults with Parkinson's, dementia, hypertension, stroke, those at pre- or post-operative care





Older adults





Systematic reviews of studies with older adults with mental health concerns



The Arts in Psychotherapy

Volume 76, November 2021, 101842



Research Article


A systematic review of music therapy interventions used to address emotional needs of older adults

Sekyung Jang a 🙎 🖾 , Lizzy Kunde b



Version published: 03 February 2017 Version history https://doi.org/10.1002/14651858.CD011022.pub2 3

Cochrane Database of Systematic Reviews



Music-based therapeutic interventions for people with dementia (Review)

van der Steen JT, Smaling HJA, van der Wouden JC, Bruinsma MS, Scholten RJPM, Vink AC

Cochrane Library

Trusted evidence. Informed decisions. Better health.

Cochrane Reviews ▼ Trials ▼ Clinical Answers ▼ About ▼ He

Cochrane Database of Systematic Reviews | Review - Intervention

Dance movement therapy for dementia

Vicky Karkou, Bonnie Meekums Authors' declarations of interest

There are neither large scale multi-national trials in arts therapies nor studies on combined approaches.



Example from research in care homes

THE LANCET Healthy Longevity

Sub

This journal Journals Publish Clinical Global health Multimedia Events About

ARTICLES | VOLUME 3, ISSUE 3, E153-E165, MARCH 2022




PDF [602 KB]

Clinical effectiveness of music interventions for dementia and depression in elderly care (MIDDEL): Australian cohort of an international pragmatic cluster-randomised controlled trial

Felicity A Baker, PhD 🔌 🖾 • Young-Eun C Lee, PhD • Tanara Vieira Sousa, PhD •

Phoebe A Stretton-Smith, MMusThrp • Jeanette Tamplin, PhD • Vigdis Sveinsdottir, PhD • et al. Show all authors

Open Access • Published: March, 2022 • DOI: https://doi.org/10.1016/S2666-7568(22)00027-7 •







The Arts and Health Office at WHO and the new WHO Jameel Arts and Health Lab



Health Topics ~

Countries ~

Newsroom ∨

Emergencies ~

Data >


About WHO >

Home / Initiatives / Arts and health





Commissioned work on the first WHO-Lancet Global Series





What supports therapeutic change?


Therapeutic factors



NYU – World Health Organization Arts and Health Launch Event, September 2021



Therapeutic factors





19 domains

1.	Embodiment (n=14)



Three domains with therapeutic factors labelled joint &/or specific

Embodiment

- Body Awareness
- Physicality of the arts
- Experiencing the body

Concretization

- Creating something visible
- Re-enactment
- Self display
- Visual narratives

Symbolism and metaphor

- Symbolic work
- Metaphor
- Unconscious processing through the arts



Neuroscience Meets Dance in Therapy

Towards mechanisms of change





























Neuroscience Meets Dance in Therapy

Neuroscience Meets Dance in Therapy

Neuroscience, Dance, the Arts and Therapy:

Translating Research into Practice

Edge Hill University, Ormskirk, UK 7th June 2024, 10:00 - 18:00

Organised by:



Dr Valentina Cazzato




Prof Vicky Karkou



Dr Stergios Makris

Dr Asaf Bachrach (CNRS France) Keynote Speaker













Humanities Research Council















Future research

- Large good quality outcome studies with manualised interventions for specific populations
- Neuroscientific studies to understand the process:
 - Participant observation
 - Synchrony and mirroring
 - Movement meditation
- Implementation studies are also urgently needed






Implementation studies

Arts4Us: a collaboration with over 50 partners

£2.5 million to scale up place-based arts initiatives that support the mental health of children and young







people





